

Physical Properties of Low-Luminosity Galaxies at $z \sim 2$

M. Lemoine-Busserolle, T. Contini, R. Pelló, J.-F. Le Borgne, & J.-P. Kneib

LAOMP - UMR 5572, 14 avenue E. Belin, F-31400 Toulouse, France

C. Lidman

European Southern Observatory, Alonso de Cordova, 3107 Vitacura, Chile

Abstract. We report the results obtained from ISAAC/VLT near-IR spectroscopy of two low-luminosity $z \sim 1.9$ galaxies located in the core of the lensing cluster AC114. The amplification factor allowed to obtain, for the first time, physical properties (SFR, abundance ratios, mass, etc) of star-forming galaxies, 1 to 2 magnitudes fainter than in previous studies of LBGs at $z \sim 3$.

Near-infrared (NIR) spectroscopy provides a view of the physical properties of the high-redshift (up to $z \sim 6$) star-forming galaxies using various rest-frame optical line diagnostics (Balmer lines, [O II] $\lambda\lambda 3726, 3728$ [O III] $\lambda\lambda 4959, 5007$ [N II] $\lambda 6584$, etc) commonly applied for nearby galaxy studies. With the recent advent of NIR spectrographs on 8-10m class telescopes, these studies have started in the $1.8 \leq z \leq 3.5$ domain (Pettini et al. 1998, 2001; Kobulnicky & Koo 2000). The main applications have been the determination of the star formation rate (SFR) deduced from the $H\alpha$ luminosity, the abundance of oxygen derived from the strong oxygen emission lines, and virial masses obtained from the emission line widths. The main goal of these studies is to derive the physical properties of distant galaxies using the same parameter space as for galaxies in the local Universe.

We report, in this contribution, the results obtained on two lensed low-luminosity $z \sim 1.9$ galaxies (S2 and A2) located in the core of the lensing cluster AC114. The amplification factor allowed to obtain, for the first time, emission-line measurements on star-forming galaxies as faint as $M_B \sim -20$, thus 1 to 2 magnitudes fainter than in previous studies of LBGs at $z \sim 3$ (e.g. Pettini et al. 2001). ISAAC/VLT medium-resolution *JHK* spectra of the two galaxies S2 and A2 have been obtained in two hour exposure (per band and per object) on 26-27 september 2001 and reduced in a standard way. Bright rest-frame optical emission lines are measured (intensity and line width) with a high S/N ratio (see Figure 1) and have been used to perform a detailed analysis of the physical properties (SFR, chemical abundances, virial mass, etc) of S2 and A2. The SFRs derived from $H\alpha$ luminosity are 20 (S2) and 40 (A2) times higher than those derived from the UV (1500Å) luminosity, without dust extinction correction. The difference reduces to a factor of 3 (S2) and 6 (A2) assuming an extinction $E(B-V)=0.3$ mags, but still the $SFR(H\alpha) > SFR(1500\text{\AA})$. These results suggest that quite large dust extinction corrections are needed for intrinsically faint optically-selected galaxies compared to other samples of bright




Figure 1. *Left:* ISAAC/VLT spectrum of the low-luminosity ~ 1.9 galaxy S2 in the core of the lensing cluster AC114. *Right:* N/O versus O/H relation for nearby (HII galaxies and SBNGs) and high- z star-forming galaxies. The location of the two low-luminosity $z \sim 1.9$ galaxies S2 and A2 is shown without extinction correction (empty squares) and assuming $E(B-V)=0.3$ (filled squares).

UV-selected distant galaxies (e.g. Pettini et al. 2001). The large spectral coverage obtained, from [O II] $\lambda\lambda 3726, 3728$ to $H\alpha$, [N II] $\lambda 6584$ allowed to set strong constraints on the abundances ratios of these two low-luminosity galaxies (see Figure 1). The behavior of S2 and A2 in terms of metallicity is very different, and they are also different from typical LBGs at $z \sim 3$. S2 is a low-metallicity object ($Z \sim 0.03 Z_{\odot}$) with a low N/O ratio, similar to those derived in the most metal-poor nearby HII galaxies. On the contrary, A2 is a metal-rich galaxy ($Z \sim 1.5 Z_{\odot}$) with a high N/O abundance ratio, similar to those derived in the most metal-rich massive SBNGs. The position of MS 1512-cB58, a lensed luminous LBG (Teplitz et al. 2000), is intermediate between these two extremes showing abundance ratios typical of low-mass SBNGs and intermediate-redshift galaxies. These results suggest different star formation history for distant galaxies of different luminosities. Larger samples of faint high- z galaxies are urgently needed to conclude, in order to set strong constraints on the galaxy evolution scenario.

Acknowledgments. Part of this work was supported by a “Conseil Régional de la Martinique” grant and the french CNRS.

References

- Contini, T., Treyer, M.-A., Sullivan, M., Ellis, R.S. 2002, MNRAS 330, 75
- Kobulnicky, H. A., Koo, D. C. 2000, ApJ 545, 712
- Kobulnicky, H. A., Zaritsky, D. 1999, ApJ 511, 118
- Pettini, M., Kellogg, M., Steidel, C. C., et al. 1998, ApJ 508, 539
- Pettini, M., Shapley, A. E., Steidel, C. C., et al. 2001, ApJ 554, 981
- Teplitz, H. I., McLean, I.S., Becklin, E.E., et al. 2000, ApJ, 533 L65